

**PPMI NX PI-2620 Tau Imaging Substudy**  
**PI-2620 Imaging**

**Note: Women of childbearing potential must have a negative urine pregnancy test result prior to the imaging scan.**

A. Assessment Date: \_\_\_\_ / \_\_\_\_ / \_\_\_\_ (mm/dd/yyyy)

**Vital Signs Measured Approximately 5 - 60 Minutes Prior to Injection**

1. Was a study physician present to evaluate the participant prior to injection?

☐ Yes

☐ No

If no, please explain:

---

---

---

2. Time vital signs measured prior to injection: \_\_\_\_:\_\_\_\_ (24-hour clock)

To be taken after participant has been supine for 1-3 minutes:

3. Supine blood pressure: \_\_\_\_ / \_\_\_\_ mmHg (systolic/diastolic)

4. Supine heart rate: \_\_\_\_ beats per minute

5. Time of 18F-PI-2620 injection: \_\_\_\_:\_\_\_\_ (24-hour clock)

**Vital Signs Measured Approximately 15 – 30 Minutes Post-Injection**

6. Time vital signs measured after injection: \_\_\_\_:\_\_\_\_ (24-hour clock)

To be taken after participant has been supine for 1-3 minutes:

7. Supine blood pressure: \_\_\_\_ / \_\_\_\_ mmHg (systolic/diastolic)

8. Supine heart rate: \_\_\_\_ beats per minute

9. Was PI-2620 NeuroEXPLORER imaging scan completed?

☐ Yes

☐ No

If no, please explain:

---

---

---

10. Was PI-2620 Conventional PET imaging scan completed?

☐ Yes

☐ No

If no, please explain:

---

---

---

11. Was a study physician (or designee) present to evaluate the participant prior to discharge?

☐ Yes

☐ No

If no, please explain:

---

---

---